

Density-functional theory for excited states: An ensemble perspective


Emmanuel Fromager

*Laboratoire de Chimie Quantique, Institut de Chimie de Strasbourg
Université de Strasbourg, Strasbourg, France.*

Outline

- (Time-independent) **DFT for excited states**: why and how?
- Connecting Kohn-Sham **orbital** energies to **real** (ground- and excited-state) energies.
- Individual exchange-correlation **functionals for excited states** (within an ensemble).


DFT and excited states



Hohenberg-Kohn theorem:

Ground- and *excited-state* energies are in principle functionals of the ground-state density n_{Ψ_0} .

(Linear response) time-dependent DFT



Practical limitations:

- **Single-reference** perturbation theory (not adequate for **nearly-degenerate** situations).
- Memory effects are absent from standard functionals (**adiabatic approximation**).
- **Multiple-electron excitations** are absent from the adiabatic TD-DFT spectrum.

(Excited) state-specific DFT

$\Psi \perp \Psi_0$




$$E_1 \equiv E_1^{\perp \Psi_0}[n_{\Psi_1}]$$

$$E_0 \text{ ————— } \Psi_0$$

M. Levy and A. Nagy, *Phys. Rev. Lett.* **83**, 4361 (1999).

In *Coulomb systems*, individual densities are sufficient, i.e.

$$E_I \equiv E^{Coul}[n_{\Psi_I}]$$



$$v(\mathbf{r}) = - \sum_A \frac{Z_A}{|\mathbf{r} - \mathbf{R}_A|}$$

P. W. Ayers, M. Levy, and A. Nagy, *Phys. Rev. A* **85**, 042518 (2012).

P. W. Ayers, M. Levy, and A. Nagy, *J. Chem. Phys.* **143**, 191101 (2015).

P. W. Ayers, M. Levy, and A. Nagy, *Theor. Chem. Acc.* **137**, 152 (2018).

DFT for (canonical) ensembles



Gross-Oliveira-Kohn (GOK) DFT:

The *ensemble energy* $E^{\mathbf{w}} := \sum_{I \geq 0} w_I E_I$ is a functional of the *ensemble density* $n^{\mathbf{w}}(\mathbf{r}) := \sum_{I \geq 0} w_I n_{\Psi_I}(\mathbf{r})$.

From DFT to GOK-DFT

Ensemble Hxc functional:

I am here to say that n is an ensemble density!

$$E_{\text{Hxc}}[n] \longrightarrow E_{\text{Hxc}}^{\mathbf{w}}[n]$$

$$\text{Example: } \varphi_{1s}^{\mathbf{w}}(\mathbf{r}) = \sqrt{(1-\mathbf{w})\varphi_{1s}^2(\mathbf{r}) + \mathbf{w}\varphi_{2s}^2(\mathbf{r})}$$

Ensemble Kohn-Sham equations:

$$\left[-\frac{\nabla^2}{2} + v_{\text{ext}}(\mathbf{r}) + \frac{\delta E_{\text{Hxc}}^{\mathbf{w}}[n]}{\delta n(\mathbf{r})} \right]_{n=n^{\mathbf{w}}} \varphi_i(\mathbf{r}) = \varepsilon_i^{\mathbf{w}} \varphi_i(\mathbf{r})$$

$$n^{\mathbf{w}}(\mathbf{r}) = \sum_{I \geq 0} \mathbf{w}_I \sum_{i \in I} |\varphi_i(\mathbf{r})|^2$$

Levy-Zahariev *shift**:

$$\varepsilon_i^{\mathbf{w}} \rightarrow \bar{\varepsilon}_i^{\mathbf{w}} = \varepsilon_i^{\mathbf{w}} + \frac{E_{\text{Hxc}}^{\mathbf{w}}[n] - \int d\mathbf{r} \frac{\delta E_{\text{Hxc}}^{\mathbf{w}}[n]}{\delta n(\mathbf{r})} n(\mathbf{r})}{\int d\mathbf{r} n(\mathbf{r})} \Bigg|_{n=n^{\mathbf{w}}}$$

*M. Levy and F. Zahariev, Phys. Rev. Lett. **113**, 113002 (2014).

Extracting individual properties from GOK-DFT

$$E_c^{\mathbf{w}}[n^{\mathbf{w}}] \equiv \sum_{I \geq 0} w_I \left(\langle \Psi_I | \hat{H} | \Psi_I \rangle - \langle \Phi_I^{\mathbf{w}} | \hat{H} | \Phi_I^{\mathbf{w}} \rangle \right)$$

Ensemble correlation energy

Ensemble KS determinant

Exact ensemble density constraint:

$$\sum_{I \geq 0} w_I n_{\Psi_I}(\mathbf{r}) = n^{\mathbf{w}}(\mathbf{r}) = \sum_{I \geq 0} w_I n_{\Phi_I^{\mathbf{w}}}(\mathbf{r})$$

How can we extract E_I and n_{Ψ_I} from the KS ensemble?

Extracting individual energies

$$\bar{\mathcal{E}}_I^{\mathbf{w}} + \sum_{J>0} (\delta_{IJ} - w_J) \left. \frac{\partial E_{\text{Hxc}}^{\mathbf{w}}[n]}{\partial w_J} \right|_{n=n^{\mathbf{w}}} = E_I$$

True energies


$$\bar{\mathcal{E}}_I^{\mathbf{w}} = \sum_{i \in I} \bar{\varepsilon}_i^{\mathbf{w}}$$

Reproduce the exact ensemble energy only


$$\mathcal{E}_I^{\mathbf{w}} = \sum_{i \in I} \varepsilon_i^{\mathbf{w}}$$

KS energies


(Two-electron) Hubbard dimer model



$$t = \frac{1}{2}, U = 1, \Delta v_{\text{ext}} = 1, w_2 = 0$$



$$t = \frac{1}{2}, U = 1, \Delta v_{\text{ext}} = 1, w_2 = \frac{1}{4}$$



Correlation functionals for ensembles: Where to start?

Individual correlations:

$$E_c^w[n^w] = \sum_{I \geq 0} w_I \langle \Psi_I | \hat{H} | \Psi_I \rangle - \sum_{I \geq 0} w_I \langle \Phi_I^w | \hat{H} | \Phi_I^w \rangle$$

$$\sum_{I \geq 0} w_I n_{\Psi_I}(\mathbf{r}) = \sum_{I \geq 0} w_I n_{\Phi_I^w}(\mathbf{r})$$

$$\sum_{I \geq 0} w_I E_{c,I}^w \equiv \sum_{I \geq 0} w_I \left(\langle \Psi_I | \hat{T} + \hat{W}_{ee} | \Psi_I \rangle - \langle \Phi_I^w | \hat{T} + \hat{W}_{ee} | \Phi_I^w \rangle \right)$$

$$n_{\Psi_I}(\mathbf{r}) \neq n_{\Phi_I^w}(\mathbf{r})$$

I am the correlation energy of state I
within the ensemble

$$E_{c,I}^w \neq \langle \Psi_I | \hat{H} | \Psi_I \rangle - \langle \Phi_I^w | \hat{H} | \Phi_I^w \rangle$$

State- and density-driven correlations in ensembles

$$E_{c,I}^w = \langle \Psi_I | \hat{T} + \hat{W}_{ee} | \Psi_I \rangle - \langle \Phi_I^w | \hat{T} + \hat{W}_{ee} | \Phi_I^w \rangle$$

$$= \langle \Psi_I | \hat{T} + \hat{W}_{ee} | \Psi_I \rangle - \langle \bar{\Phi}_I | \hat{T} + \hat{W}_{ee} | \bar{\Phi}_I \rangle$$

State-specific
KS determinant

$n_{\Psi_I}(\mathbf{r}) = n_{\bar{\Phi}_I}(\mathbf{r})$

I am the *state-driven* correlation energy


$$= \langle \Psi_I | \hat{H} | \Psi_I \rangle - \langle \bar{\Phi}_I | \hat{H} | \bar{\Phi}_I \rangle$$

$$+ \langle \bar{\Phi}_I | \hat{T} + \hat{W}_{ee} | \bar{\Phi}_I \rangle - \langle \Phi_I^w | \hat{T} + \hat{W}_{ee} | \Phi_I^w \rangle$$


I am the *density-driven* correlation energy

Non-uniqueness or -existence of state-driven KS states

Non-interacting Hubbard dimer



Interacting Hubbard dimer



[Skip to main content](#)

[arXiv.org](#) > [physics](#) > arXiv:2001.08605

[Download PDF](#)

Physics > Chemical Physics

Title: Individual correlations in ensemble density-functional theory: State-driven/density-driven decomposition without additional Kohn-Sham systems

Authors: [Emmanuel Fromager](#)

(Submitted on 23 Jan 2020 ([v1](#)), last revised 28 Jan 2020 (this version, v2))

Abstract: Gould and Pittalis [Phys. Rev. Lett. 123, 016401 (2019)] recently revealed a density-driven correlation energy in many-electron ensembles that must be accounted for by approximations. We show that referring to auxiliary state-driven Kohn-Sham (KS) systems, which was inherent to its evaluation, is in fact not needed. Instead, individual-state densities can be extracted directly from the KS ensemble. On that basis, a simpler and more general expression is derived and tested. The importance of density-driven effects is thus confirmed, and a direct route to approximations is introduced.

State-/density-driven decomposition without additional KS systems

- There is *no need* to introduce additional KS wave functions:

$$n_{\Psi_I}(\mathbf{r}) = n^{\mathbf{w}}(\mathbf{r}) + \sum_{J>0} (\delta_{IJ} - w_J) \frac{\partial n^{\mathbf{w}}(\mathbf{r})}{\partial w_J} \longrightarrow \text{Can be obtained from a static ensemble Casida-like equation}$$

- Exact expressions* for **state-driven** (SD) and **density-driven** (DD) correlation energies:


$$E_{c,I}^{\text{SD}} := E_c^{\mathbf{w}}[n^{\mathbf{w}}] + \sum_{J>0} (\delta_{IJ} - w_J) \frac{dE_c^{\mathbf{w}}[n^{\mathbf{w}}]}{dw_J}$$

$$E_{c,I}^{\text{DD}} := \int d\mathbf{r} \frac{\delta E_c^{\mathbf{w}}[n^{\mathbf{w}}]}{\delta n(\mathbf{r})} \left(n_{\Phi_I^{\mathbf{w}}}(\mathbf{r}) - n_{\Psi_I}(\mathbf{r}) \right)$$

Application to the Hubbard dimer

$$\frac{\Delta v_{\text{ext}}}{t} \ll \ll \frac{t}{U} \ll \ll 1$$

First (singlet) correlation energy per unit of $U(U\Delta v_{\text{ext}})^2/(4t^4)$




Conclusions and perspectives

- *Individual* energies and densities can be extracted *exactly* from GOK-DFT.
- A general and exact *SD/DD decomposition* has been derived.
- The approach is applicable to *grand canonical* ensembles (gaps, quantum embedding,...).
- Local SD correlation functionals can be extracted from *finite* uniform electron gases:
collaboration with Pierre-François Loos (Toulouse).
- Connections with *imaginary TD-DFT* [$t \rightarrow -i\tau$] under investigation.
- Extraction of *(non-adiabatic) couplings* from GOK-DFT under investigation.


References:

- K. Deur and E. Fromager, *J. Chem. Phys.* **150**, 094106 (2019).
E. Fromager, *arXiv:2001.08605* (2020).
B. Senjean and E. Fromager, *Phys. Rev. A* **98**, 022513 (2018).
B. Senjean and E. Fromager, *arXiv:1912.07125* (2020).

Funding and acknowledgements



LABEX, University of Strasbourg



Laurent Mazouin, E.F., Killian Deur, and *Bruno Senjean (now in Leiden)*

N-centered grand canonical ensembles

density:

$$n^{\mathcal{N}}(\mathbf{r}) = (1-\alpha)n^N(\mathbf{r}) + \alpha n^{N-1}(\mathbf{r})$$




electron number:

$$\mathcal{N} = N - \alpha$$

energy:

$$E(\mathcal{N}) = (1-\alpha)E^N + \alpha E^{N-1}$$



density:

$$n^{\{N,\alpha\}}(\mathbf{r}) = (1-\alpha)n^N(\mathbf{r}) + \frac{N\alpha}{N-1}n^{N-1}(\mathbf{r})$$



electron number:


$$N$$

energy:

$$\mathcal{E}^{\{N,\alpha\}} = (1-\alpha)E^N + \frac{N\alpha}{N-1}E^{N-1}$$

$$\left[\left(1 - \frac{\alpha}{N}\right) - \frac{\alpha(1-\alpha)}{N} \frac{\partial}{\partial \alpha} \right]$$

Real picture



N-centered picture