


Quantum embedding in electronic structure theory

Part 3: Exact embedding of localised orbitals for non-interacting electrons

Emmanuel Fromager


*Laboratoire de Chimie Quantique, Institut de Chimie de Strasbourg,
Université de Strasbourg, Strasbourg, France.*

Non-interacting delocalised representation



$$\hat{H} \equiv \sum_{PQ} \langle \varphi_P | \hat{h} | \varphi_Q \rangle \hat{a}_P^\dagger \hat{a}_Q$$

Non-interacting (delocalised) molecular orbital representation




$$\hat{H} \equiv \sum_{PQ} \langle \varphi_P | \hat{h} | \varphi_Q \rangle \hat{a}_P^\dagger \hat{a}_Q$$

The molecular spin-orbitals are simply obtained
by solving the **one-electron Schrödinger equation**

$$\hat{h}\varphi_Q(\mathbf{x}) = \varepsilon_Q \varphi_Q(\mathbf{x})$$


Non-interacting (delocalised) molecular orbital representation

$\varepsilon_{\mathcal{M}}$	—	$\varphi_{\mathcal{M}}$
$\varepsilon_{\mathcal{M}-1}$	—	$\varphi_{\mathcal{M}-1}$
	⋮	
ε_{N+4}	—	φ_{N+4}
ε_{N+3}	—	φ_{N+3}
ε_{N+2}	—	φ_{N+2}
ε_{N+1}	—	φ_{N+1}
	↔	
ε_N	—	φ_N
	—	
	—	
	⋮	
ε_2	—	φ_2
ε_1	—	φ_1



$$\varphi_P(\mathbf{x}) = \sum_{\nu} C_{\nu P} \chi_{\nu}(\mathbf{x})$$

Non-interacting (delocalised) molecular orbital representation




$$\hat{H} \equiv \sum_{PQ} \langle \varphi_P | \hat{h} | \varphi_Q \rangle \hat{a}_P^\dagger \hat{a}_Q$$

The molecular spin-orbitals are simply obtained
by solving the **one-electron Schrödinger equation**

$$\hat{h}\varphi_Q(\mathbf{x}) = \varepsilon_Q \varphi_Q(\mathbf{x})$$


$$\langle \varphi_P | \hat{h} | \varphi_Q \rangle = \varepsilon_P \delta_{PQ}$$

Non-interacting (delocalised) molecular orbital representation



$$\begin{aligned}\hat{H} &\equiv \sum_{PQ} \langle \varphi_P | \hat{h} | \varphi_Q \rangle \hat{a}_P^\dagger \hat{a}_Q \\ &= \sum_P \varepsilon_P \hat{a}_P^\dagger \hat{a}_P\end{aligned}$$


Non-interacting (delocalised) molecular orbital representation



$$\begin{aligned}\hat{H} &\equiv \sum_{PQ} \langle \varphi_P | \hat{h} | \varphi_Q \rangle \hat{a}_P^\dagger \hat{a}_Q \\ &= \sum_P \varepsilon_P \hat{a}_P^\dagger \hat{a}_P\end{aligned}$$

The exact solutions to the non-interacting Schrödinger equation
are Slater determinants $\hat{a}_{P_1}^\dagger \hat{a}_{P_2}^\dagger \dots \hat{a}_{P_{N-1}}^\dagger \hat{a}_{P_N}^\dagger | \text{vac} \rangle$

Non-interacting (delocalised) molecular orbital representation




$$\begin{aligned}\hat{H} &\equiv \sum_{PQ} \langle \varphi_P | \hat{h} | \varphi_Q \rangle \hat{a}_P^\dagger \hat{a}_Q \\ &= \sum_P \varepsilon_P \hat{a}_P^\dagger \hat{a}_P\end{aligned}$$

The exact solutions to the non-interacting Schrödinger equation
are Slater determinants $\hat{a}_{P_1}^\dagger \hat{a}_{P_2}^\dagger \dots \hat{a}_{P_{N-1}}^\dagger \hat{a}_{P_N}^\dagger | \text{vac} \rangle$

$$\hat{H} \hat{a}_{P_1}^\dagger \hat{a}_{P_2}^\dagger \dots \hat{a}_{P_{N-1}}^\dagger \hat{a}_{P_N}^\dagger | \text{vac} \rangle = \left(\sum_{i=1}^N \varepsilon_{P_i} \right) \hat{a}_{P_1}^\dagger \hat{a}_{P_2}^\dagger \dots \hat{a}_{P_{N-1}}^\dagger \hat{a}_{P_N}^\dagger | \text{vac} \rangle$$

Non-interacting (delocalised) molecular orbital representation




$$\hat{H} = \sum_P \varepsilon_P \hat{a}_P^\dagger \hat{a}_P$$

Spin-orbital occupation operator


$$\hat{H} \hat{a}_{P_1}^\dagger \hat{a}_{P_2}^\dagger \dots \hat{a}_{P_{N-1}}^\dagger \hat{a}_{P_N}^\dagger | \text{vac} \rangle = \left(\sum_{i=1}^N \varepsilon_{P_i} \right) \hat{a}_{P_1}^\dagger \hat{a}_{P_2}^\dagger \dots \hat{a}_{P_{N-1}}^\dagger \hat{a}_{P_N}^\dagger | \text{vac} \rangle$$

Non-interacting (delocalised) molecular orbital representation




$$|\Psi_0\rangle = \hat{a}_1^\dagger \hat{a}_2^\dagger \dots \hat{a}_{N-1}^\dagger \hat{a}_N^\dagger |\text{vac}\rangle$$

1RDM in the molecular orbital (mo) representation



$$\gamma_{PQ}^{mo} = \langle \Psi_0 | \hat{a}_P^\dagger \hat{a}_Q | \Psi_0 \rangle$$



$$| \Psi_0 \rangle = \hat{a}_1^\dagger \hat{a}_2^\dagger \dots \hat{a}_{N-1}^\dagger \hat{a}_N^\dagger | \text{vac} \rangle$$

1RDM in the molecular orbital (mo) representation

$$\gamma_{PQ}^{mo} = \langle \Psi_0 | \hat{a}_P^\dagger \hat{a}_Q | \Psi_0 \rangle \quad \text{where} \quad |\Psi_0\rangle = \hat{a}_1^\dagger \hat{a}_2^\dagger \dots \hat{a}_{N-1}^\dagger \hat{a}_N^\dagger |vac\rangle$$


1RDM in the molecular orbital (mo) representation

$$\gamma_{PQ}^{mo} = \langle \Psi_0 | \hat{a}_P^\dagger \hat{a}_Q | \Psi_0 \rangle \quad \text{where} \quad |\Psi_0\rangle = \hat{a}_1^\dagger \hat{a}_2^\dagger \dots \hat{a}_{N-1}^\dagger \hat{a}_N^\dagger |vac\rangle$$


Is φ_Q occupied in Ψ_0 ?

1RDM in the molecular orbital (mo) representation

$$\gamma_{PQ}^{mo} = \langle \Psi_0 | \hat{a}_P^\dagger \hat{a}_Q | \Psi_0 \rangle \quad \text{where} \quad |\Psi_0\rangle = \hat{a}_1^\dagger \hat{a}_2^\dagger \dots \hat{a}_{N-1}^\dagger \hat{a}_N^\dagger |vac\rangle$$

Is φ_Q occupied in Ψ_0 ?


NO

0

1RDM in the molecular orbital (mo) representation

$$\gamma_{PQ}^{mo} = \langle \Psi_0 | \hat{a}_P^\dagger \hat{a}_Q | \Psi_0 \rangle \quad \text{where} \quad |\Psi_0\rangle = \hat{a}_1^\dagger \hat{a}_2^\dagger \dots \hat{a}_{N-1}^\dagger \hat{a}_N^\dagger |vac\rangle$$

↔




1RDM in the molecular orbital (mo) representation

$$\gamma_{PQ}^{mo} = \langle \Psi_0 | \hat{a}_P^\dagger \hat{a}_Q | \Psi_0 \rangle \quad \text{where} \quad |\Psi_0\rangle = \hat{a}_1^\dagger \hat{a}_2^\dagger \dots \hat{a}_{N-1}^\dagger \hat{a}_N^\dagger |vac\rangle$$

↔

Is φ_Q occupied in Ψ_0 ?




If φ_P is occupied in Ψ_0 then $\hat{a}_P^\dagger \hat{a}_Q | \Psi_0 \rangle = 0$ **Pauli principle**

1RDM in the molecular orbital (mo) representation

$$\gamma_{PQ}^{mo} = \langle \Psi_0 | \hat{a}_P^\dagger \hat{a}_Q | \Psi_0 \rangle \quad \text{where} \quad |\Psi_0\rangle = \hat{a}_1^\dagger \hat{a}_2^\dagger \dots \hat{a}_{N-1}^\dagger \hat{a}_N^\dagger |vac\rangle$$

↔




If φ_P is unoccupied in Ψ_0 then $\hat{a}_P^\dagger \hat{a}_Q | \Psi_0 \rangle \perp | \Psi_0 \rangle \Rightarrow \langle \Psi_0 | \hat{a}_P^\dagger \hat{a}_Q | \Psi_0 \rangle = 0$

1RDM in the molecular orbital (mo) representation


$$\gamma_{PQ}^{mo} = \langle \Psi_0 | \hat{a}_P^\dagger \hat{a}_Q | \Psi_0 \rangle \quad \text{where} \quad |\Psi_0\rangle = \hat{a}_1^\dagger \hat{a}_2^\dagger \dots \hat{a}_{N-1}^\dagger \hat{a}_N^\dagger |vac\rangle$$

↔



1RDM in the molecular orbital (mo) representation

$$\gamma_{PQ}^{mo} = \langle \Psi_0 | \hat{a}_P^\dagger \hat{a}_Q | \Psi_0 \rangle \quad \text{where} \quad |\Psi_0\rangle = \hat{a}_1^\dagger \hat{a}_2^\dagger \dots \hat{a}_{N-1}^\dagger \hat{a}_N^\dagger |vac\rangle$$





1RDM in the molecular orbital (mo) representation

$$\gamma_{PQ}^{mo} = \langle \Psi_0 | \hat{a}_P^\dagger \hat{a}_Q | \Psi_0 \rangle$$

where


$$| \Psi_0 \rangle = \hat{a}_1^\dagger \hat{a}_2^\dagger \dots \hat{a}_{N-1}^\dagger \hat{a}_N^\dagger | \text{vac} \rangle$$





1RDM in the molecular orbital (mo) representation

$$\gamma_{PQ}^{mo} = \langle \Psi_0 | \hat{a}_P^\dagger \hat{a}_Q | \Psi_0 \rangle \quad \text{where} \quad |\Psi_0\rangle = \hat{a}_1^\dagger \hat{a}_2^\dagger \dots \hat{a}_{N-1}^\dagger \hat{a}_N^\dagger |vac\rangle$$

↔



1RDM in the molecular orbital (mo) representation





$$\gamma^{mo} = \begin{bmatrix} \varphi_1 & \varphi_2 & \dots & \varphi_N & \dots & \varphi_Q & \dots & \varphi_{\mathcal{M}} \\ \varphi_1 & 1 & 1 & 1 & \dots & 0 & \dots & 0 \\ \varphi_2 & 1 & 1 & 1 & \dots & 1 & \dots & 1 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots \\ \varphi_P & 0 & 0 & 0 & \dots & 1 & \dots & 1 \\ \varphi_N & 0 & 0 & 0 & \dots & 1 & \dots & 1 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \ddots & \vdots \\ \varphi_{\mathcal{M}} & 0 & 0 & 0 & \dots & 0 & \dots & 0 \end{bmatrix}$$

Non-interacting problem solved!



1RDM in the molecular orbital (mo) representation



$$\gamma^{mo} = \begin{bmatrix} \varphi_1 & \varphi_2 & \dots & \varphi_N & \dots & \varphi_Q & \dots & \varphi_{\mathcal{M}} \end{bmatrix}$$

$$\begin{bmatrix} \varphi_1 \\ \varphi_2 \\ \vdots \\ \varphi_P \\ \varphi_N \\ \vdots \\ \varphi_{\mathcal{M}} \end{bmatrix}$$

The matrix γ^{mo} is a diagonal matrix where the diagonal elements are binary values (0 or 1). The diagonal starts at 1 for φ_1 , goes down to 0 for φ_Q , and then continues with 1s for the remaining orbitals up to $\varphi_{\mathcal{M}}$. The off-diagonal elements are all 0.

No entanglement between the molecular orbitals
in the non-interacting case

Idempotency property

$$\gamma^{mo} = \begin{matrix} & \varphi_1 \varphi_2 & \dots & \varphi_N & \dots & \varphi_Q & \dots & \varphi_M \\ \varphi_1 & 1 & 1 & 1 & & & & \\ \varphi_2 & 1 & 0 & & & & & \\ \varphi_P & 0 & & 1 & & & & \\ \varphi_N & & \dots & 1 & 1 & & & \\ & & & & & 0 & & \\ & & & & & & 0 & \\ \varphi_M & & & & & & & 0 \end{matrix} = [\gamma^{mo}]^2$$

Turning to the localised picture (useless here although interesting)

$$|\varphi_P\rangle = \sum_I C_{IP} |\chi_I\rangle$$

Localised spin-orbitals

$$|\varphi_Q\rangle = \sum_J C_{JQ} |\chi_J\rangle$$

Turning to the localised picture (useless here although interesting)

$$|\varphi_P\rangle = \sum_I C_{IP} |\chi_I\rangle$$

Localised spin-orbitals

$$|\varphi_Q\rangle = \sum_J C_{JQ} |\chi_J\rangle$$

$$\langle\varphi_P|\varphi_Q\rangle = \delta_{PQ} = \sum_{IJ} C_{IP} C_{JQ} \langle\chi_I|\chi_J\rangle$$

Turning to the localised picture (useless here although interesting)

$$|\varphi_P\rangle = \sum_I C_{IP} |\chi_I\rangle$$
$$|\varphi_Q\rangle = \sum_J C_{JQ} |\chi_J\rangle$$

Localised spin-orbitals

$$\langle \varphi_P | \varphi_Q \rangle = \delta_{PQ} = \sum_{IJ} C_{IP} C_{JQ} \langle \chi_I | \chi_J \rangle$$

Orthonormalisation procedure

$$\delta_{IJ}$$

Turning to the localised picture (useless here although interesting)

$$|\varphi_P\rangle = \sum_I C_{IP} |\chi_I\rangle$$
$$|\varphi_Q\rangle = \sum_J C_{JQ} |\chi_J\rangle$$

Localised spin-orbitals

*Molecular orbital
coefficients matrix*

$$\langle\varphi_P|\varphi_Q\rangle = \delta_{PQ} = \sum_{IJ} C_{IP} C_{JQ} \delta_{IJ}$$
$$= \sum_I C_{IP} C_{IQ} = \sum_I [\mathbf{C}^T]_{PI} [\mathbf{C}]_{IQ}$$
$$= [\mathbf{C}^T \mathbf{C}]_{PQ}$$

Turning to the localised picture (useless here although interesting)

$$|\varphi_P\rangle = \sum_I C_{IP} |\chi_I\rangle$$
$$|\varphi_Q\rangle = \sum_J C_{JQ} |\chi_J\rangle$$

Localised spin-orbitals

*Unitary transformation
from the delocalised to localised pictures*

$$\mathbf{C}^{-1} = \mathbf{C}^T$$

$$\delta_{PQ} = [\mathbf{C}^T \mathbf{C}]_{PQ}$$

Turning to the localised picture (useless here although interesting)

$$\gamma_{PQ}^{mo} = \langle \Psi_0 | \hat{a}_P^\dagger \hat{a}_Q | \Psi_0 \rangle$$



$$\begin{aligned}\hat{a}_P^\dagger &= \sum_I C_{IP} \hat{c}_I^\dagger \\ \hat{a}_Q &= \sum_J C_{JQ} \hat{c}_J\end{aligned}$$

$$\gamma_{PQ}^{mo} = \sum_{IJ} C_{IP} C_{JQ} \langle \Psi_0 | \hat{c}_I^\dagger \hat{c}_J | \Psi_0 \rangle$$

Turning to the localised picture (useless here although interesting)

$$\gamma_{PQ}^{mo} = \langle \Psi_0 | \hat{a}_P^\dagger \hat{a}_Q | \Psi_0 \rangle$$

$$\begin{aligned}\hat{a}_P^\dagger &= \sum_I C_{IP} \hat{c}_I^\dagger \\ \hat{a}_Q &= \sum_J C_{JQ} \hat{c}_J\end{aligned}$$



1RDM in the
delocalised representation

$$\gamma_{PQ}^{mo} = \sum_{IJ} C_{IP} C_{JQ} \boxed{\langle \Psi_0 | \hat{c}_I^\dagger \hat{c}_J | \Psi_0 \rangle}$$

$$\gamma_{IJ}^{loc} \equiv \gamma_{IJ}$$

Turning to the localised picture (useless here although interesting)

$$\gamma_{PQ}^{mo} = \langle \Psi_0 | \hat{a}_P^\dagger \hat{a}_Q | \Psi_0 \rangle$$



$$\begin{aligned}\hat{a}_P^\dagger &= \sum_I C_{IP} \hat{c}_I^\dagger \\ \hat{a}_Q &= \sum_J C_{JQ} \hat{c}_J\end{aligned}$$

$$\gamma_{PQ}^{mo} = \sum_{IJ} C_{IP} C_{JQ} \gamma_{IJ} = [\mathbf{C}^T]_{PI} \gamma_{IJ}^{loc} [\mathbf{C}]_{JQ}$$

Turning to the localised picture (useless here although interesting)

$$\gamma_{PQ}^{mo} = \langle \Psi_0 | \hat{a}_P^\dagger \hat{a}_Q | \Psi_0 \rangle$$



$$\begin{aligned}\hat{a}_P^\dagger &= \sum_I C_{IP} \hat{c}_I^\dagger \\ \hat{a}_Q &= \sum_J C_{JQ} \hat{c}_J\end{aligned}$$

$$\begin{aligned}\gamma_{PQ}^{mo} &= \sum_{IJ} C_{IP} C_{JQ} \gamma_{IJ} = [\mathbf{C}^T]_{PI} \gamma_{IJ}^{loc} [\mathbf{C}]_{JQ} \\ &= [\mathbf{C}^T \gamma^{loc} \mathbf{C}]_{PQ}\end{aligned}$$

Turning to the localised picture (useless here although interesting)

$$\gamma^{loc} = \mathbf{C}\gamma^{mo}\mathbf{C}^T$$

$$\gamma^{mo} = \mathbf{C}^T\gamma^{loc}\mathbf{C}$$

$$\gamma_{PQ}^{mo} = [\mathbf{C}^T\gamma^{loc}\mathbf{C}]_{PQ}$$

Turning to the localised picture (useless here although interesting)

$$\gamma^{loc} = \mathbf{C}\gamma^{mo}\mathbf{C}^T$$

$$\gamma_{IJ}^{loc} = \sum_{PQ} C_{IP} \gamma_{PQ}^{mo} C_{JQ} = \sum_P \text{occupied spin-MOs} C_{IP} C_{JP}$$

Turning to the localised picture (useless here although interesting)

$$\gamma^{loc} = \mathbf{C}\gamma^{mo}\mathbf{C}^T \quad \xleftarrow{\text{Not diagonal!}}$$

$$\gamma_{IJ}^{loc} = \langle \hat{c}_I^\dagger \hat{c}_J \rangle_{\Psi_0} = \sum_P \text{occupied spin-MOs} C_{IP} C_{JP} \neq \delta_{IJ}$$

Turning to the localised picture (useless here although interesting)

$$\gamma^{loc} = \mathbf{C}\gamma^{mo}\mathbf{C}^T \quad \xleftarrow{\text{Not diagonal!}}$$

$$\gamma_{IJ}^{loc} = \langle \hat{c}_I^\dagger \hat{c}_J \rangle_{\Psi_0} = \sum_P \text{occupied spin-MOs} C_{IP} C_{JP} \neq \delta_{IJ}$$

Any localised spin-orbital χ_I is **entangled** with the other spin-orbitals χ_J

Turning to the localised picture (useless here although interesting)


$$\gamma^{loc} = \mathbf{C}\gamma^{mo}\mathbf{C}^T \quad \xleftarrow{\text{Not diagonal!}}$$

$$\gamma_{IJ}^{loc} = \langle \hat{c}_I^\dagger \hat{c}_J \rangle_{\Psi_0} = \sum_P \text{occupied spin-MOs} C_{IP} C_{JP} \neq \delta_{IJ}$$


Any localised spin-orbital χ_I is **entangled** with the other spin-orbitals χ_J

unlike in the delocalised molecular orbital space!


Prototypical ring of $L = 16$ hydrogen atoms




Prototypical ring of $L = 16$ hydrogen atoms




Prototypical ring of $L = 16$ hydrogen atoms




To-be-embedded (so-called *impurity*)
localised orbital



Exact density matrix functional embedding




Two-electron system




N -electron system

Exact density matrix functional embedding



Two-electron system

Let's prove that the embedding is **exact** for **non-interacting** electrons




N -electron system

The “Householder embedding” project




From left to right: **S. Sekaran** (Strasbourg, France),
M. Saubanère (Montpellier, France),
L. Mazouin (Strasbourg, France), and E.F.




E.F. and M. Tsuchiizu (Nara, Japan).


The Householder transformation



The Householder transformation




The Householder transformation




The Householder transformation

Normalized
Householder vector


$$\frac{\mathbf{X} - \mathbf{Y}}{|\mathbf{X} - \mathbf{Y}|} = \mathbf{v}$$




The Householder transformation




The Householder transformation




The Householder transformation




The Householder transformation




The Householder transformation




The Householder transformation



The Householder transformation



The Householder transformation




The Householder transformation

$$P^{-1} = P^T = P$$


Unitary transformation

$$P = I - 2vv^T$$


The Householder transformation




Householder transformed density matrix embedding




Householder transformed density matrix embedding




Householder transformed density matrix embedding



Householder transformed density matrix embedding




Householder transformed density matrix embedding




Householder transformed density matrix embedding


$$|\mathbf{X}|^2 - \gamma_{00}^2 = |\mathbf{Y}|^2 - \gamma_{00}^2 = \tilde{\gamma}_{10}^2 = \sum_{J>0} \gamma_{J0}^2$$



Householder transformed density matrix embedding




Householder transformed density matrix embedding



Householder transformed density matrix embedding

$$\mathbf{v} = \frac{\mathbf{X} - \mathbf{Y}}{|\mathbf{X} - \mathbf{Y}|} \quad \xleftarrow{\text{Householder vector}}$$

where $|\mathbf{X} - \mathbf{Y}|^2 = 2 \left(|\mathbf{Y}|^2 - \mathbf{X}^T \mathbf{Y} \right) = 2 (\tilde{\gamma}_{10}^2 - \gamma_{10} \tilde{\gamma}_{10}) = 2 \tilde{\gamma}_{10} (\tilde{\gamma}_{10} - \gamma_{10})$




Householder transformed density matrix embedding

$$\mathbf{v} = \frac{\mathbf{X} - \mathbf{Y}}{|\mathbf{X} - \mathbf{Y}|} \quad \text{Householder vector}$$

where

$$|\mathbf{X} - \mathbf{Y}|^2 = \pm 2 \sqrt{\sum_{J>0} \gamma_{J0}^2} \left(\pm \sqrt{\sum_{J>0} \gamma_{J0}^2} - \gamma_{10} \right)$$




If one single neighbour...

$$|\mathbf{X} - \mathbf{Y}|^2 = \pm 2 |\gamma_{10}| (\pm |\gamma_{10}| - \gamma_{10})$$

Householder transformed density matrix embedding

$$\mathbf{v} = \frac{\mathbf{X} - \mathbf{Y}}{|\mathbf{X} - \mathbf{Y}|} \quad \text{Householder vector}$$

where


$$|\mathbf{X} - \mathbf{Y}|^2 = \pm 2 \sqrt{\sum_{J>0} \gamma_{J0}^2} \left(\pm \sqrt{\sum_{J>0} \gamma_{J0}^2} - \gamma_{10} \right)$$


If one single neighbour...


$$|\mathbf{X} - \mathbf{Y}|^2 = \pm 2 |\gamma_{10}| (\pm |\gamma_{10}| - \gamma_{10})$$

choose $-\operatorname{sgn}(\gamma_{10}) \leftarrow |\mathbf{X} - \mathbf{Y}| \neq 0$


Householder transformed density matrix embedding



Householder transformed density matrix embedding



Householder transformed density matrix embedding



The Householder transformation is an **explicit functional** of the density matrix!

Householder representation in second quantization

$$\mathbf{P} \equiv \mathbf{P}[\gamma] = \mathbf{I} - 2\mathbf{v}\mathbf{v}^T$$

Unitary Householder transformation matrix

Householder representation in second quantization

$$\mathbf{P} \equiv \mathbf{P}[\gamma] = \mathbf{I} - 2\mathbf{v}\mathbf{v}^T$$

Unitary Householder transformation matrix

$$P_{IJ} = \delta_{IJ} - 2v_I v_J$$

*Householder transformation **matrix elements***

Householder representation in second quantization

$$\mathbf{P} \equiv \mathbf{P}[\gamma] = \mathbf{I} - 2\mathbf{v}\mathbf{v}^T \quad \textit{Unitary Householder transformation matrix}$$

$$P_{IJ} = \delta_{IJ} - 2v_I v_J \quad \textit{Householder transformation matrix elements}$$

$$\hat{d}_I^\dagger = \sum_J P_{IJ} \hat{c}_J^\dagger \quad \textit{Creates delocalised Householder orbitals}$$

Householder representation in second quantization

$$\hat{d}_I^\dagger = \sum_J P_{IJ} \hat{c}_J^\dagger$$

*From the **localised** to the **Householder** representation*

Householder representation in second quantization

$$\hat{d}_I^\dagger = \sum_J P_{IJ} \hat{c}_J^\dagger$$

*From the **localised** to the **Householder** representation*

$$\sum_I P_{KI} \hat{d}_I^\dagger = \sum_J \sum_I P_{KI} P_{IJ} \hat{c}_J^\dagger$$

Householder representation in second quantization

$$\hat{d}_I^\dagger = \sum_J P_{IJ} \hat{c}_J^\dagger$$

*From the **localised** to the **Householder** representation*

$$\sum_I P_{KI} \hat{d}_I^\dagger = \sum_J \sum_I P_{KI} P_{IJ} \hat{c}_J^\dagger = \sum_J [\mathbf{P}^2]_{KJ} \hat{c}_J^\dagger = \sum_J \delta_{KJ} \hat{c}_J^\dagger = \hat{c}_K^\dagger$$

Householder representation in second quantization

$$\hat{d}_I^\dagger = \sum_J P_{IJ} \hat{c}_J^\dagger$$

*From the **localised** to the **Householder** representation*

$$\hat{c}_K^\dagger = \sum_I P_{KI} \hat{d}_I^\dagger$$

*From the **Householder** to the **localised** representation*

Householder representation in second quantization


$$\mathbf{v} = \frac{\mathbf{X} - \mathbf{Y}}{|\mathbf{X} - \mathbf{Y}|}$$

Householder vector

$$\hat{d}_0^\dagger = \sum_J P_{0J} \hat{c}_J^\dagger \quad v_0 = 0 \quad \hat{c}_0^\dagger$$

$\underbrace{}$

$$\delta_{0J} - 2v_0 v_J$$



The **impurity is invariant** under the transformation

Householder representation in second quantization

$$\mathbf{P} \equiv \mathbf{P}[\gamma] = \mathbf{I} - 2\mathbf{v}\mathbf{v}^T$$

Unitary Householder transformation matrix

$$P_{IJ} = \delta_{IJ} - 2v_I v_J$$

Householder transformation matrix elements

$$\hat{d}_I^\dagger = \sum_J P_{IJ} \hat{c}_J^\dagger$$

*Creates delocalised **Householder orbitals***

$$\hat{d}_0^\dagger = \hat{c}_0^\dagger$$

*The **impurity** is **invariant** under the transformation*

$$\hat{d}_1^\dagger |\text{vac}\rangle = \sum_J P_{1J} |\chi_J\rangle$$

*Will play the role of the **bath orbital***

1RDM in the Householder representation

$$\tilde{\gamma}_{J0} = \langle \Psi_0 | \hat{d}_J^\dagger \hat{d}_0 | \Psi_0 \rangle$$


$$= \sum_I P_{JI} \langle \Psi_0 | \hat{c}_I^\dagger \hat{c}_0 | \Psi_0 \rangle$$

$$= \sum_I P_{JI} \gamma_{I0}$$

$$= [\mathbf{P}\mathbf{X}]_J$$

$$= [\mathbf{Y}]_J$$

$$\stackrel{J \geq 1}{=} 0$$



1RDM in the Householder representation

$$\tilde{\gamma}_{J0} = \langle \Psi_0 | \hat{d}_J^\dagger \hat{d}_0 | \Psi_0 \rangle$$


$$= \sum_I P_{JI} \langle \Psi_0 | \hat{c}_I^\dagger \hat{c}_0 | \Psi_0 \rangle$$

$$= \sum_I P_{JI} \gamma_{I0}$$

$$= [\mathbf{P}\mathbf{X}]_J$$

$$= [\mathbf{Y}]_J$$

$$\stackrel{J \geq 1}{=} 0$$



By construction, the *impurity* is **entangled only with the bath**

1RDM in the Householder representation

Theorem: As the electrons are **non-interacting**, the **bath** turns out to be **entangled only with the impurity**

1RDM in the Householder representation

Theorem: As the electrons are **non-interacting**, the **bath** turns out to be **entangled only with the impurity**

Proof

$$\tilde{\gamma} = \tilde{\gamma}^2 \quad \text{Idempotency property}$$

1RDM in the Householder representation

Theorem: As the electrons are **non-interacting**, the **bath** turns out to be **entangled only with the impurity**

Proof

$$\tilde{\gamma} = \tilde{\gamma}^2 \quad \text{Idempotency property}$$

$$\begin{aligned}\tilde{\gamma}_{J0} &= \langle \Psi_0 | \hat{d}_J^\dagger \hat{d}_0 | \Psi_0 \rangle = [\tilde{\gamma}^2]_{J0} = \sum_K \tilde{\gamma}_{JK} \tilde{\gamma}_{K0} \\ &= \tilde{\gamma}_{J0} \tilde{\gamma}_{00} + \tilde{\gamma}_{J1} \tilde{\gamma}_{10} + \sum_{K>1} \tilde{\gamma}_{JK} \times 0\end{aligned}$$

1RDM in the Householder representation

Theorem: As the electrons are **non-interacting**, the **bath** turns out to be **entangled only with the impurity**

Proof

$$\tilde{\gamma} = \tilde{\gamma}^2 \quad \text{Idempotency property}$$

$$\tilde{\gamma}_{J0} = \tilde{\gamma}_{J0}\tilde{\gamma}_{00} + \tilde{\gamma}_{J1}\tilde{\gamma}_{10}$$



$$\tilde{\gamma}_{J1} = \frac{\tilde{\gamma}_{J0} (1 - \tilde{\gamma}_{00})}{\tilde{\gamma}_{10}}$$

1RDM in the Householder representation

Theorem: As the electrons are **non-interacting**, the **bath** turns out to be **entangled only with the impurity**

Proof

$$\tilde{\gamma}_{J1} = \frac{\tilde{\gamma}_{J0} (1 - \tilde{\gamma}_{00})}{\tilde{\gamma}_{10}}$$

$$J > 1 \quad \tilde{\gamma}_{J0} = 0$$

No entanglement between the **impurity** and the orbitals other than the bath



$$\tilde{\gamma}_{J1} = 0$$

1RDM in the Householder representation

Theorem: As the electrons are **non-interacting**, the **bath** turns out to be **entangled only with the impurity**

Proof

No entanglement between the **bath** and the orbitals other than the **impurity!**




$$\tilde{\gamma}_{J1} = \langle \Psi_0 | \hat{d}_J^\dagger \hat{d}_1 | \Psi_0 \rangle = 0$$

1RDM in the Householder representation

Theorem: As the electrons are **non-interacting**, the **bath** turns out to be **entangled only with the impurity**

Proof

$$\tilde{\gamma}_{J1} = \frac{\tilde{\gamma}_{J0} (1 - \tilde{\gamma}_{00})}{\tilde{\gamma}_{10}}$$



$$\tilde{\gamma}_{11} + \tilde{\gamma}_{00} = 1$$

1RDM in the Householder representation

Theorem: As the electrons are **non-interacting**, the **bath** turns out to be **entangled only with the impurity**

Proof

$$\tilde{\gamma}_{11} + \tilde{\gamma}_{00} = \langle \Psi_0 | \hat{d}_1^\dagger \hat{d}_1 | \Psi_0 \rangle + \langle \Psi_0 | \hat{d}_0^\dagger \hat{d}_0 | \Psi_0 \rangle = 1$$

The “**impurity+bath**” cluster contains exactly **one electron (per spin)**

1RDM in the Householder representation

Theorem: As the electrons are **non-interacting**, the **bath** turns out to be **entangled only with the impurity**


Proof

$$\tilde{\gamma}_{11} + \tilde{\gamma}_{00} = \langle \Psi_0 | \hat{d}_1^\dagger \hat{d}_1 | \Psi_0 \rangle + \langle \Psi_0 | \hat{d}_0^\dagger \hat{d}_0 | \Psi_0 \rangle = 1$$

The “**impurity+bath**” cluster contains exactly **one electron (per spin)**

The cluster is a **closed quantum system** that can be described with a two-electron wave function Ψ^C


1RDM in the Householder representation



Energy evaluation by fragmentation

$$\hat{H} = \sum_{IJ} \bar{h}_{IJ} \hat{c}_I^\dagger \hat{c}_J \quad \textcolor{magenta}{\textit{Localised representation}}$$


Energy evaluation by fragmentation



$$\hat{H} = \sum_{IJ} \bar{h}_{IJ} \hat{c}_I^\dagger \hat{c}_J$$

Localised representation

Energy evaluation by fragmentation




$$\hat{H} = \sum_{IJ} \bar{h}_{IJ} \hat{c}_I^\dagger \hat{c}_J$$

Localised representation

Energy contributions involving the impurity

$$2 \sum_J \bar{h}_{0J} \langle \Psi_0 | \hat{c}_0^\dagger \hat{c}_J | \Psi_0 \rangle$$

Energy evaluation by fragmentation



$$\hat{H} = \sum_{IJ} \bar{h}_{IJ} \hat{c}_I^\dagger \hat{c}_J$$

Localised representation


Energy contributions involving the impurity

$$2 \sum_J \bar{h}_{0J} \langle \Psi_0 | \hat{c}_0^\dagger \hat{c}_J | \Psi_0 \rangle$$

Householder representation \longrightarrow $= 2 \sum_K \left(\sum_J \bar{h}_{0J} P_{JK} \right) \langle \Psi_0 | \hat{d}_0^\dagger \hat{d}_K | \Psi_0 \rangle$

\tilde{h}_{0K}

Energy evaluation by fragmentation



$$\hat{H} = \sum_{IJ} \bar{h}_{IJ} \hat{c}_I^\dagger \hat{c}_J$$

Localised representation


Energy contributions involving the impurity

$$2 \sum_J \bar{h}_{0J} \langle \Psi_0 | \hat{c}_0^\dagger \hat{c}_J | \Psi_0 \rangle$$

$$= 2 \sum_K \left(\sum_J \bar{h}_{0J} P_{JK} \right) \langle \Psi_0 | \hat{d}_0^\dagger \hat{d}_K | \Psi_0 \rangle$$

$$= 2 \sum_K \tilde{h}_{0K} \tilde{\gamma}_{0K} = 2 \left(\tilde{h}_{00} \tilde{\gamma}_{00} + \tilde{h}_{01} \tilde{\gamma}_{01} \right)$$

Energy evaluation by fragmentation



Determined from the cluster

$$2 \sum_J \bar{h}_{0J} \langle \Psi_0 | \hat{c}_0^\dagger \hat{c}_J | \Psi_0 \rangle = 2 \left(\tilde{h}_{00} \tilde{\gamma}_{00} + \tilde{h}_{01} \tilde{\gamma}_{01} \right)$$

Approximate embedding for interacting electrons

The present embedding approach is useless for non-interacting electrons (!)

Approximate embedding for interacting electrons

The present embedding approach is *useless for non-interacting electrons* (!)

We need $\gamma_{IJ} = \gamma_{IJ}^{loc} = \sum_P^{\text{occupied spin-MOs}} C_{IP} C_{JP}$

Approximate embedding for interacting electrons

We need

$$\gamma_{IJ} = \gamma_{IJ}^{loc} = \sum_P \text{occupied spin-MOs} C_{IP} C_{JP}$$

We have to solve the Schrödinger equation for the **full system!**

Approximate embedding for interacting electrons

*The present embedding approach is **useless for non-interacting electrons** (!)*

*However, it can be used for describing electron **repulsions** (within the cluster)*

Approximate embedding for interacting electrons

The present embedding approach is useless for non-interacting electrons (!)

However, it can be used for describing electron repulsions (within the cluster)

$$\hat{H} \equiv \sum_{PQ} \tilde{h}_{PQ} \hat{d}_P^\dagger \hat{d}_Q + \frac{1}{2} \sum_{PQRS} \tilde{g}_{PQRS} \hat{d}_P^\dagger \hat{d}_Q^\dagger \hat{d}_S \hat{d}_R$$

Full Hamiltonian in the Householder representation

Approximate embedding for interacting electrons

The present embedding approach is *useless for non-interacting electrons* (!)

However, it can be used for describing electron *repulsions (within the cluster)*

$$\hat{H} \equiv \sum_{PQ} \tilde{h}_{PQ} \hat{d}_P^\dagger \hat{d}_Q + \frac{1}{2} \sum_{PQRS} \tilde{g}_{PQRS} \hat{d}_P^\dagger \hat{d}_Q^\dagger \hat{d}_S \hat{d}_R$$




Projection onto the cluster


$$\hat{H}^{\mathcal{C}} \equiv \sum_{P,Q \in \mathcal{C}} \tilde{h}_{PQ} \hat{d}_P^\dagger \hat{d}_Q + \frac{1}{2} \sum_{P,Q,R,S \in \mathcal{C}} \tilde{g}_{PQRS} \hat{d}_P^\dagger \hat{d}_Q^\dagger \hat{d}_S \hat{d}_R$$

Application to the 1D Hubbard model

The “Householder embedding” project




From left to right: **S. Sekaran** (Strasbourg, France),
M. Saubanère (Montpellier, France),
L. Mazouin (Strasbourg, France), and E.F.




E.F. and M. Tsuchiizu (Nara, Japan).

Prototypical ring of L hydrogen atoms




Prototypical ring of L hydrogen atoms



Hubbard model

$$\hat{H} \approx -t \sum_{\sigma=\uparrow,\downarrow} \sum_{i=0}^{L-1} \left(\hat{c}_{i\sigma}^\dagger \hat{c}_{(i+1)\sigma} + \hat{c}_{(i+1)\sigma}^\dagger \hat{c}_{i\sigma} \right) + U \sum_{i=0}^{L-1} \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\downarrow}^\dagger \hat{c}_{i\downarrow} \hat{c}_{i\uparrow}$$

Prototypical ring of L hydrogen atoms




Hubbard model

$$\hat{H} \approx -t \sum_{\sigma=\uparrow,\downarrow} \sum_{i=0}^{L-1} \left(\hat{c}_{i\sigma}^\dagger \hat{c}_{(i+1)\sigma} + \hat{c}_{(i+1)\sigma}^\dagger \hat{c}_{i\sigma} \right) + U \sum_{i=0}^{L-1} \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\downarrow}^\dagger \hat{c}_{i\downarrow} \hat{c}_{i\uparrow}$$

Hückel model $-t \equiv \beta$

Prototypical ring of L hydrogen atoms



Hubbard model

$$\hat{H} \approx -t \sum_{\sigma=\uparrow,\downarrow} \sum_{i=0}^{L-1} \left(\hat{c}_{i\sigma}^\dagger \hat{c}_{(i+1)\sigma} + \hat{c}_{(i+1)\sigma}^\dagger \hat{c}_{i\sigma} \right) + U \sum_{i=0}^{L-1} \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\downarrow}^\dagger \hat{c}_{i\downarrow} \hat{c}_{i\uparrow}$$

Two-electron repulsion
on each atom only

Prototypical ring of L hydrogen atoms

$$U/t \ll 1$$


Weakly correlated regime

$$U/t \gg 1$$


Strongly correlated regime

Hubbard model

$$\hat{H} \approx -t \sum_{\sigma=\uparrow,\downarrow} \sum_{i=0}^{L-1} \left(\hat{c}_{i\sigma}^\dagger \hat{c}_{(i+1)\sigma} + \hat{c}_{(i+1)\sigma}^\dagger \hat{c}_{i\sigma} \right) + U \sum_{i=0}^{L-1} \hat{c}_{i\uparrow}^\dagger \hat{c}_{i\downarrow}^\dagger \hat{c}_{i\downarrow} \hat{c}_{i\uparrow}$$



$L = 400$ atoms



Ht-DMFET per-site energies away from half-filling ($n < 1$)

