

Introduction to second quantization

Emmanuel Fromager



Institut de Chimie de Strasbourg - Laboratoire de Chimie Quantique -
Université de Strasbourg /CNRS

ISTPC 2015 summer school, Aussois, France, june 2015

Electronic Hamiltonian in first quantization

N -electron Hamiltonian within the Born-Oppenheimer approximation:

$$\hat{H} = \hat{T} + \hat{V}_{\text{ne}} + \hat{W}_{\text{ee}}$$

$$\hat{T} = \sum_{i=1}^N \hat{t}(i) \quad \text{where} \quad \hat{t}(i) \equiv -\frac{1}{2} \nabla_{\mathbf{r}_i}^2 \quad \rightarrow \quad \text{kinetic energy}$$

$$\hat{V}_{\text{ne}} = \sum_{i=1}^N \hat{v}_{\text{ne}}(i) \quad \text{where} \quad \hat{v}_{\text{ne}}(i) \equiv - \sum_A^{\text{nuclei}} \frac{Z_A}{|\mathbf{r}_i - \mathbf{R}_A|} \times \quad \rightarrow \quad \text{electron-nuclei attraction}$$

$$\hat{W}_{\text{ee}} = \frac{1}{2} \sum_{i \neq j}^N \hat{w}_{\text{ee}}(i, j) \quad \text{where} \quad \hat{w}_{\text{ee}}(i, j) \equiv \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|} \times \quad \rightarrow \quad \text{electron-electron repulsion}$$

One-electron wavefunction

- Let us start with **Schrödinger theory**: the quantum state of a single electron is written as

$$|\Psi\rangle = \int d\mathbf{r} \Psi(\mathbf{r})|\mathbf{r}\rangle$$

where $\Psi(\mathbf{r})$ is the one-electron wavefunction (**orbital**) and $|\mathbf{r}\rangle$ denotes the quantum state "the electron is at position \mathbf{r} ". In other words, $\hat{\mathbf{r}}|\mathbf{r}\rangle = \mathbf{r}|\mathbf{r}\rangle$.

- This choice of basis is known as "**r representation**".
- Orthonormalization** condition: $\langle \mathbf{r}' | \mathbf{r} \rangle = \delta(\mathbf{r}' - \mathbf{r})$ \leftarrow Dirac distribution

Useful formulas: (1) " $\forall f$ ", $\int d\mathbf{r} f(\mathbf{r})\delta(\mathbf{r}' - \mathbf{r}) = f(\mathbf{r}')$

$$(2) \quad \delta(\mathbf{r}' - \mathbf{r}) = \frac{1}{(2\pi)^3} \int d\mathbf{k} e^{i\mathbf{k} \cdot (\mathbf{r}' - \mathbf{r})}$$

- In this representation, the **momentum** vector operator is defined as follows,

$$\hat{\mathbf{p}}|\Psi\rangle = -i \int d\mathbf{r} \nabla_{\mathbf{r}} \Psi(\mathbf{r})|\mathbf{r}\rangle$$

- "**k representation**":

$$|\mathbf{k}\rangle = \frac{1}{(2\pi)^{3/2}} \int d\mathbf{r} e^{i\mathbf{k}\cdot\mathbf{r}}|\mathbf{r}\rangle$$

Note that $\hat{\mathbf{p}}|\mathbf{k}\rangle = \mathbf{k}|\mathbf{k}\rangle$ and $\langle\mathbf{k}'|\mathbf{k}\rangle = \delta(\mathbf{k}' - \mathbf{k})$ \longleftarrow use formula (2) !

- Pauli theory**: the **spin** of the electron is now considered as an additional degree of freedom. The quantum state of a single electron is then written as

$$|\Psi\rangle = \int d\mathbf{r} \sum_{\sigma=\alpha,\beta} \Psi(\mathbf{r}, \sigma)|\mathbf{r}, \sigma\rangle$$

where $|\mathbf{r}, \alpha\rangle$ denotes the quantum state "electron at position \mathbf{r} with **spin up**" and $|\mathbf{r}, \beta\rangle$ corresponds to the state "electron at position \mathbf{r} with **spin down**"

Two-electron wavefunction

- In the **non-relativistic** case, a single electron will have a spin σ_0 which is either up or down. The corresponding wavefunction Ψ^{σ_0} can then be written as a **spin-orbital** $\Psi^{\sigma_0}(\mathbf{r}, \sigma) = \Psi(\mathbf{r})\delta_{\sigma\sigma_0}$.
- With the notations $X = (\mathbf{r}, \sigma)$ and $\int dX = \int d\mathbf{r} \sum_{\sigma=\alpha,\beta}$,

a one-electron quantum state in Pauli theory is simply written as

$$|\Psi\rangle = \int dX \Psi(X) |X\rangle$$

- **Two-electron** case:

$$|\Psi\rangle = \int \int dX_1 dX_2 \Psi(X_1, X_2) |\mathbf{1}: X_1, \mathbf{2}: X_2\rangle$$

where the two-electron quantum state $|\mathbf{1}: X_1, \mathbf{2}: X_2\rangle$ corresponds to "**electron 1** in state $|X_1\rangle$ and **electron 2** in state $|X_2\rangle$ "

- **Anti-symmetrization** principle: a physical two-electron wavefunction should fulfill the condition

$$\boxed{\Psi(X_1, X_2) = -\Psi(X_2, X_1)}$$

thus leading to

$$|\Psi\rangle = - \int \int dX_1 dX_2 \Psi(X_2, X_1) |\textcolor{red}{1}: \boxed{X_1}, \textcolor{blue}{2}: X_2\rangle = - \underbrace{\int \int dX_1 dX_2 \Psi(X_1, X_2) |\textcolor{red}{1}: X_2, \textcolor{blue}{2}: \boxed{X_1}\rangle}_{|\Psi_{\textcolor{red}{1}\leftrightarrow\textcolor{blue}{2}}\rangle}$$

and

$$|\Psi\rangle = \frac{1}{2} \int \int dX_1 dX_2 \underbrace{\left[\Psi(X_1, X_2) - \Psi(X_2, X_1) \right]}_{0 \text{ if } X_1 = X_2} |\textcolor{red}{1}: X_1, \textcolor{blue}{2}: X_2\rangle$$

Conclusion: the anti-symmetrization of the wavefunction ensures that electrons are **indistinguishable** and that they cannot be in the same quantum state (**Pauli principle**).

Slater determinants

- Let $\{\varphi_K(X)\}_K$ denote an **orthonormal** basis of (molecular) spin-orbitals. Two electrons that occupy the spin-orbitals $\varphi_I(X)$ and $\varphi_J(X)$ will be described by the (normalized) **Slater determinant**

$$\Phi_{IJ}(X_1, X_2) = \frac{1}{\sqrt{2}} \begin{vmatrix} \varphi_I(X_1) & \varphi_I(X_2) \\ \varphi_J(X_1) & \varphi_J(X_2) \end{vmatrix} = \frac{1}{\sqrt{2}} \left(\varphi_I(X_1)\varphi_J(X_2) - \varphi_I(X_2)\varphi_J(X_1) \right)$$

- Note that Slater determinants and, consequently, linear combinations of Slater determinants are **anti-symmetric**.
- Therefore, Slater determinants are convenient "**building blocks**" for computing the electronic wavefunction.
- Still, we may wonder if we really need this **complicated** expression obtained from the determinant (obviously things get worse for a larger number of electrons).
- Another drawback of the current formulation: both Slater determinant and Hamiltonian expressions **depend on the number of electrons**

"What is occupied ?" rather than "Who occupies what ?"

- Since electrons are **indistinguishable**, there is no need to know that electron 1 occupies φ_I and electron 2 occupies φ_J or the other way around ...
- The important information is that spin-orbitals φ_I and φ_J are **occupied** and the remaining ones are **empty**.
- **Second quantization** is a formalism that relies on this idea.
- At the beginning, there was "nothing" ... $|\text{vac}\rangle$ \longleftarrow normalized "**vacuum** state"
- ... then was introduced the concept of **annihilation** of an electron occupying φ_I , that would obviously give zero when applied to the vacuum state: $\boxed{\forall I, \hat{a}_I |\text{vac}\rangle = 0}$ **(rule 1)**
- ... and then came the concept of **creation** of an electron occupying φ_I : $\hat{a}_I^\dagger |\text{vac}\rangle \equiv |\varphi_I\rangle$

"What is occupied ?" rather than "Who occupies what ?"

- ... and then came the idea to create another electron occupying φ_J : $\hat{a}_J^\dagger \hat{a}_I^\dagger |\text{vac}\rangle \equiv |\Phi_{IJ}\rangle$
- Note that the creation operator \hat{a}_I^\dagger is the **adjoint** of the annihilation operator \hat{a}_I . This ensures, in particular, that one-electron and vacuum states are orthogonal:

$$\langle \varphi_I | \text{vac} \rangle = \langle \hat{a}_I^\dagger \text{vac} | \text{vac} \rangle = \langle \text{vac} | \hat{a}_I | \text{vac} \rangle = 0$$

- In order to have a representation that is equivalent to the one used in first quantization, we only need two more rules:

$$\boxed{\forall I, J, \quad [\hat{a}_I, \hat{a}_J]_+ = \hat{a}_I \hat{a}_J + \hat{a}_J \hat{a}_I = 0} \quad \text{(rule 2)} \quad \longrightarrow \quad [\hat{a}_I^\dagger, \hat{a}_J^\dagger]_+ = [\hat{a}_J, \hat{a}_I]_+^\dagger = 0$$

$$\boxed{\forall I, J, \quad [\hat{a}_I, \hat{a}_J^\dagger]_+ = \hat{a}_I \hat{a}_J^\dagger + \hat{a}_J^\dagger \hat{a}_I = \delta_{IJ}} \quad \text{(rule 3)}$$

"What is occupied ?" rather than "Who occupies what ?"

- **Rule 2** contains the indistinguishability of the electrons, $\hat{a}_J^\dagger \hat{a}_I^\dagger |\text{vac}\rangle = -\hat{a}_I^\dagger \hat{a}_J^\dagger |\text{vac}\rangle$,

and the Pauli principle, $\hat{a}_I^\dagger \hat{a}_I^\dagger |\text{vac}\rangle = 0$.

- **Rule 3** ensures that you can only annihilate what has already been created (!),

$$\hat{a}_I \hat{a}_J^\dagger |\text{vac}\rangle = \delta_{IJ} |\text{vac}\rangle - \hat{a}_J^\dagger \hat{a}_I |\text{vac}\rangle = \delta_{IJ} |\text{vac}\rangle$$

- It is now very easy to generate representations of Slater determinants for an **arbitrary number N of electrons**: multiply more creation operators !

$$|I_1 I_2 \dots I_{N-1} I_N\rangle = \hat{a}_{I_1}^\dagger \hat{a}_{I_2}^\dagger \dots \hat{a}_{I_{N-1}}^\dagger \hat{a}_{I_N}^\dagger |\text{vac}\rangle \quad \equiv \quad \frac{1}{\sqrt{N}} \det [\varphi_{I_i}(X_j)]$$

EXERCISE: (1) Show that $|I_1 I_2 \dots I_{N-1} I_N\rangle$ is **normalized**.

(2) Let us consider another state $|J_1 J_2 \dots J_{N-1} J_N\rangle$ and assume that at least one of the occupied spin-orbitals (let us denote it φ_{J_k}) is not occupied in $|I_1 I_2 \dots I_{N-1} I_N\rangle$. Show that the two states are **orthogonal**.

(3) The "counting" operator \hat{N} is defined as $\hat{N} = \sum_I \hat{n}_I$ where $\hat{n}_I = \hat{a}_I^\dagger \hat{a}_I$. Show that

$$\begin{aligned} \hat{n}_I |I_1 I_2 \dots I_{N-1} I_N\rangle &= |I_1 I_2 \dots I_{N-1} I_N\rangle & \text{if } I = I_k \quad 1 \leq k \leq N \\ &= 0 & \text{otherwise} \end{aligned}$$

and conclude that $\hat{N} |I_1 I_2 \dots I_{N-1} I_N\rangle = N |I_1 I_2 \dots I_{N-1} I_N\rangle$.

(4) Explain why states corresponding to different numbers of electrons are automatically orthogonal.

(5) Explain why any state $|\Psi\rangle$ fulfills the condition $0 \leq \langle \Psi | \hat{n}_I | \Psi \rangle \leq 1$.

One-electron operators in second quantization

- Let \hat{h} denote a **one-electron** operator ($\hat{t} + \hat{v}_{\text{ne}}$ for example): it acts on the one-electron states $|\varphi_I\rangle$.
- Resolution of the identity: $\sum_I |\varphi_I\rangle\langle\varphi_I| = \hat{1}$,

which leads to the **conventional representation** $\hat{h} = \hat{1} \hat{h} \hat{1} = \sum_{I,J} \langle\varphi_I|\hat{h}|\varphi_J\rangle |\varphi_I\rangle\langle\varphi_J|$.

- **Second-quantized representation:**

$$\hat{h} \equiv \sum_{I,J} \langle\varphi_I|\hat{h}|\varphi_J\rangle \hat{a}_I^\dagger \hat{a}_J$$

Indeed,

$$\begin{aligned} \left(\sum_{I,J} \langle\varphi_I|\hat{h}|\varphi_J\rangle \hat{a}_I^\dagger \hat{a}_J \right) |\varphi_K\rangle &= \left(\sum_{I,J} \langle\varphi_I|\hat{h}|\varphi_J\rangle \hat{a}_I^\dagger \hat{a}_J \right) \hat{a}_K^\dagger |\text{vac}\rangle = \left(\sum_{I,J} \langle\varphi_I|\hat{h}|\varphi_J\rangle \hat{a}_I^\dagger \delta_{JK} \right) |\text{vac}\rangle \\ &= \sum_I \langle\varphi_I|\hat{h}|\varphi_K\rangle |\varphi_I\rangle = \hat{h} |\varphi_K\rangle \end{aligned}$$

- What is convenient is that this second-quantized representation is **valid for any number N of electrons**:

$$\sum_{i=1}^N \hat{h}(i) \equiv \sum_{I,J} \langle \varphi_I | \hat{h} | \varphi_J \rangle \hat{a}_I^\dagger \hat{a}_J \equiv \hat{h}$$

The information about N has been completely transferred to the states. It **does not appear in the operator** anymore.

EXERCISE: Let us consider another orthonormal basis $\{\tilde{\varphi}_K(X)\}_K$ of spin-orbitals that we decompose in the current basis as follows, $|\tilde{\varphi}_P\rangle = \sum_Q U_{QP} |\varphi_Q\rangle$.

- (1) Show that the matrix U is unitary ($U^\dagger = U^{-1}$).
- (2) Explain why $\hat{a}_{\tilde{P}}^\dagger = \sum_Q U_{QP} \hat{a}_Q^\dagger$ and show that $\hat{h} \equiv \sum_{I,J} \langle \tilde{\varphi}_I | \hat{h} | \tilde{\varphi}_J \rangle \hat{a}_{\tilde{I}}^\dagger \hat{a}_{\tilde{J}}$.
- (3) Show how the diagonalization of \hat{h} in the one-electron space leads automatically to the diagonalization in the N -electron space (use exercise page 11).

Two-electron operators in second quantization

- Let \hat{w} denote a **two-electron** operator: it acts on two-electron states $|\varphi_I \varphi_J\rangle = |\textcolor{red}{1}: \varphi_I, \textcolor{blue}{2}: \varphi_J\rangle$.
- A **complete anti-symmetrized basis** should be used for describing the two electrons:

$$|IJ\rangle = \frac{1}{\sqrt{2}} \left(|\varphi_I \varphi_J\rangle - |\varphi_J \varphi_I\rangle \right) \equiv \hat{a}_{\textcolor{red}{I}}^\dagger \hat{a}_{\textcolor{red}{J}}^\dagger |\text{vac}\rangle \quad \text{with } I < J.$$

Consequently, any two-electron anti-symmetrized state $|\Psi\rangle$ should fulfill the condition

$$\hat{P}_{\mathcal{A}} |\Psi\rangle = |\Psi\rangle \quad \text{where} \quad \hat{P}_{\mathcal{A}} = \sum_{I < J} |IJ\rangle \langle IJ| \quad \longleftarrow \text{projection operator !}$$

- Projection of the two-electron operator onto the space of anti-symmetrized states:

$$\hat{w}_{\mathcal{A}} = \hat{P}_{\mathcal{A}} \hat{w} \hat{P}_{\mathcal{A}} = \sum_{I < J, K < L} \langle IJ | \hat{w} | KL \rangle |IJ\rangle \langle KL|$$

Two-electron operators in second quantization

EXERCISE: Prove that $\hat{w}_{\mathcal{A}} \equiv \frac{1}{2} \sum_{IJKL} \langle \varphi_I \varphi_J | \hat{w} | \varphi_K \varphi_L \rangle \hat{a}_I^\dagger \hat{a}_J^\dagger \hat{a}_L \hat{a}_K$

hint: apply $\hat{w}_{\mathcal{A}}$ and the proposed second-quantized representation to $|PQ\rangle \equiv \hat{a}_P^\dagger \hat{a}_Q^\dagger |\text{vac}\rangle$ ($P < Q$). Conclude.

- What is convenient is that this second-quantized representation is **valid for any number N of electrons** and includes the projection onto **anti-symmetrized** states:

$$\frac{1}{2} \sum_{i \neq j}^N \hat{w}(i, j) \equiv \frac{1}{2} \sum_{IJKL} \langle \varphi_I \varphi_J | \hat{w} | \varphi_K \varphi_L \rangle \hat{a}_I^\dagger \hat{a}_J^\dagger \hat{a}_L \hat{a}_K \equiv \hat{w}$$

Summary

- In summary, the electronic Hamiltonian can be written in second quantization as follows,

$$\hat{H} = \sum_{IJ} \langle \varphi_I | \hat{h} | \varphi_J \rangle \hat{a}_I^\dagger \hat{a}_J + \frac{1}{2} \sum_{IJKL} \langle \varphi_I \varphi_J | \hat{w}_{ee} | \varphi_K \varphi_L \rangle \hat{a}_I^\dagger \hat{a}_J^\dagger \hat{a}_L \hat{a}_K$$

where $\langle \varphi_I | \hat{h} | \varphi_J \rangle = \int dX \varphi_I^*(X) (\hat{h} \varphi_J)(X)$ \leftarrow one-electron integrals

$\langle \varphi_I \varphi_J | \hat{w}_{ee} | \varphi_K \varphi_L \rangle = \int \int dX_1 dX_2 \varphi_I^*(X_1) \varphi_J^*(X_2) (\hat{w}_{ee} \varphi_K \varphi_L)(X_1, X_2)$ \leftarrow two-electron integrals

- Note that this expression is also valid for a **relativistic** Hamiltonian. Two or four-component **spinors** should be used rather than spin-orbitals in conjunction with the **Dirac** (Breit) Coulomb Hamiltonian.
- The standard (**non-relativistic**) Hamiltonian will be used in the following.

EXERCISE:

(1) At the non-relativistic level, **real algebra** can be used, $\varphi_I(X) = \varphi_{i\sigma}(\mathbf{r}, \tau) = \phi_i(\mathbf{r})\delta_{\sigma\tau}$,

$$\hat{h} \equiv -\frac{1}{2}\nabla_{\mathbf{r}}^2 + v_{\text{ne}}(\mathbf{r}) \quad \text{and} \quad \hat{w}_{\text{ee}} \equiv \frac{1}{|\mathbf{r}_1 - \mathbf{r}_2|}.$$

Show that the Hamiltonian, that is here a spin-free operator, can be rewritten in the basis of the molecular orbitals $\{\phi_p(\mathbf{r})\}_p$ as follows

$$\hat{H} = \sum_{p,q} h_{pq} \hat{E}_{pq} + \frac{1}{2} \sum_{p,q,r,s} \langle pr|qs \rangle \left(\hat{E}_{pq} \hat{E}_{rs} - \delta_{qr} \hat{E}_{ps} \right)$$

where $\hat{E}_{pq} = \sum_{\sigma} \hat{a}_{p,\sigma}^{\dagger} \hat{a}_{q,\sigma}$, $h_{pq} = \langle \phi_p | \hat{h} | \phi_q \rangle$ and

$$\langle pr|qs \rangle = \int \int d\mathbf{r}_1 d\mathbf{r}_2 \phi_p(\mathbf{r}_1) \phi_r(\mathbf{r}_2) \frac{1}{|\mathbf{r}_1 - \mathbf{r}_2|} \phi_q(\mathbf{r}_1) \phi_s(\mathbf{r}_2) = (pq|rs)$$

EXERCISE:

For any normalized N -electron wavefunction Ψ , we define the one-electron (1) and two-electron (2) **reduced density matrices** (RDM) as follows,

$$D_{pq} = \langle \Psi | \hat{E}_{pq} | \Psi \rangle \quad \text{and} \quad D_{pqrs} = \langle \Psi | \hat{E}_{pq} \hat{E}_{rs} - \delta_{qr} \hat{E}_{ps} | \Psi \rangle.$$

(1) Show that the **1RDM is symmetric** and that $\forall p$, the **occupation** $n_p = D_{pp}$ of the orbital p fulfills the inequality $0 \leq n_p \leq 2$. Show that the trace of the 1RDM equals N .

(2) Explain why the expectation value for the **energy** $\langle \Psi | \hat{H} | \Psi \rangle$ can be **determined from the 2RDM**.

Hint: show that $D_{pq} = \frac{1}{N-1} \sum_r D_{pqrr}$.

(3) Let us consider the particular case $|\Psi\rangle \rightarrow |\Phi\rangle = \prod_{i=1}^{N/2} \prod_{\sigma} \hat{a}_{i,\sigma}^{\dagger} |\text{vac}\rangle$. Explain why both density matrices are non-zero only in the occupied-orbital space.

Show that $D_{ij} = 2\delta_{ij}$ and $D_{ijkl} = 4\delta_{ij}\delta_{kl} - 2\delta_{jk}\delta_{il}$ and ...

... deduce the corresponding energy expression:

$$\langle \Phi | \hat{H} | \Phi \rangle = 2 \sum_{i=1}^{N/2} h_{ii} + \sum_{i,j=1}^{N/2} \left(2 \langle ij | ij \rangle - \langle ij | ji \rangle \right).$$

(4) Let i, j and a, b denote occupied and unoccupied (virtuals) orbitals in Φ , respectively. Explain why \hat{E}_{ai} and $\hat{E}_{ai}\hat{E}_{bj}$ are referred to as **single** excitation and **double excitation operators**, respectively.

Hint: derive simplified expressions for $|\Phi_i^a\rangle = \frac{1}{\sqrt{2}}\hat{E}_{ai}|\Phi\rangle$ and $|\Phi_{ij}^{ab}\rangle = \frac{1}{2}\hat{E}_{ai}\hat{E}_{bj}|\Phi\rangle$ with $i < j, a < b$.

Why "second" quantization ?

- Let us consider a **single electron** occupying the spin-orbital $\Psi(X) = \Psi(\mathbf{r}, \sigma)$. The corresponding expectation value for the nuclear potential energy equals

$$\langle \Psi | \hat{v}_{\text{ne}} | \Psi \rangle = \int dX \Psi^*(X) v_{\text{ne}}(\mathbf{r}) \Psi(X) = \int d\mathbf{r} v_{\text{ne}}(\mathbf{r}) \underbrace{\sum_{\sigma} \Psi^*(\mathbf{r}, \sigma) \Psi(\mathbf{r}, \sigma)}_{n(\mathbf{r}): \text{ density}}$$

- For an arbitrary number of electrons: $\hat{V}_{\text{ne}} = \sum_{i=1}^N \hat{v}_{\text{ne}}(i) \equiv \sum_{p,q} \langle \phi_p | \hat{v}_{\text{ne}} | \phi_q \rangle \hat{E}_{pq}$

- Change of basis: $\{|\phi_p\rangle\}_p \longrightarrow \{|\mathbf{r}\rangle\}_{\mathbf{r} \in \mathbb{R}^3},$

$$\langle \phi_p | \hat{v}_{\text{ne}} | \phi_q \rangle \longrightarrow \langle \mathbf{r}' | \hat{v}_{\text{ne}} | \mathbf{r} \rangle = v_{\text{ne}}(\mathbf{r}) \delta(\mathbf{r}' - \mathbf{r})$$

$$\hat{a}_{p,\sigma}^\dagger \longrightarrow \hat{a}_{\mathbf{r},\sigma}^\dagger = \hat{\Psi}^\dagger(\mathbf{r}, \sigma)$$

thus leading to

$$\hat{V}_{\text{ne}} = \int d\mathbf{r} v_{\text{ne}}(\mathbf{r}) \hat{n}(\mathbf{r})$$

← **local potential** operator !

where the **density operator** equals $\hat{n}(\mathbf{r}) = \sum_{\sigma} \hat{\Psi}^{\dagger}(\mathbf{r}, \sigma) \hat{\Psi}(\mathbf{r}, \sigma)$.

- Note that the **electron density** associated with the normalized N -electron wavefunction Ψ equals $n_{\Psi}(\mathbf{r}) = \langle \Psi | \hat{n}(\mathbf{r}) | \Psi \rangle$.
- Note also that $\int d\mathbf{r} \hat{n}(\mathbf{r}) = \hat{N} \longrightarrow \int d\mathbf{r} n_{\Psi}(\mathbf{r}) = \langle \Psi | \hat{N} | \Psi \rangle = N$
- In **practice**, the density is usually obtained from the molecular orbitals and the 1RDM:

$$\begin{aligned} |\mathbf{r}, \sigma\rangle &= \sum_p |\phi_p, \sigma\rangle \langle \phi_p | \mathbf{r} \rangle = \sum_p \phi_p(\mathbf{r}) |\phi_p, \sigma\rangle &\longrightarrow \hat{n}(\mathbf{r}) &= \sum_{pq} \phi_p(\mathbf{r}) \phi_q(\mathbf{r}) \hat{E}_{pq} \\ & &\longrightarrow n_{\Psi}(\mathbf{r}) &= \sum_{pq} \phi_p(\mathbf{r}) \phi_q(\mathbf{r}) D_{pq} \end{aligned}$$

EXERCISE:

(1) Show that the kinetic energy operator is written as follows in the **k representation**:

$$\hat{T} = \sum_{i=1}^N \frac{1}{2} \widehat{\mathbf{p}^2}(i) \equiv \int d\mathbf{k} \frac{k^2}{2} \sum_{\sigma} \hat{\Psi}^{\dagger}(\mathbf{k}, \sigma) \hat{\Psi}(\mathbf{k}, \sigma) \quad \text{where} \quad \hat{\Psi}^{\dagger}(\mathbf{k}, \sigma) = \hat{a}_{\mathbf{k}, \sigma}^{\dagger}.$$

(2) Explain why $\hat{\Psi}^{\dagger}(\mathbf{k}, \sigma) = \frac{1}{(2\pi)^{3/2}} \int d\mathbf{r} e^{i\mathbf{k} \cdot \mathbf{r}} \hat{\Psi}^{\dagger}(\mathbf{r}, \sigma)$ and

$$\hat{\Psi}^{\dagger}(\mathbf{r}, \sigma) = \frac{1}{(2\pi)^{3/2}} \int d\mathbf{k} e^{-i\mathbf{k} \cdot \mathbf{r}} \hat{\Psi}^{\dagger}(\mathbf{k}, \sigma)$$

(3) Conclude that

$$\hat{T} \equiv -\frac{1}{2} \int d\mathbf{r} \sum_{\sigma} \hat{\Psi}^{\dagger}(\mathbf{r}, \sigma) \nabla_{\mathbf{r}}^2 \hat{\Psi}(\mathbf{r}, \sigma)$$

(4) Let $\hat{n}_1(\mathbf{r}', \mathbf{r}) = \sum_{\sigma} \hat{\Psi}^{\dagger}(\mathbf{r}', \sigma) \hat{\Psi}(\mathbf{r}, \sigma)$ denote the one-electron **density matrix operator**. Deduce from question (3) the standard expression

$$\hat{T} \equiv -\frac{1}{2} \int \int d\mathbf{r} d\mathbf{r}' \delta(\mathbf{r}' - \mathbf{r}) \nabla_{\mathbf{r}}^2 \hat{n}_1(\mathbf{r}', \mathbf{r})$$

(5) Show that the one-electron density matrix operator is connected with the \hat{E}_{pq} operators as follows,

$$\hat{n}_1(\mathbf{r}', \mathbf{r}) = \sum_{pq} \phi_p(\mathbf{r}') \phi_q(\mathbf{r}) \hat{E}_{pq}$$

(6) The one-electron density matrix associated with the wavefunction Ψ is defined as $n_1(\mathbf{r}', \mathbf{r}) = \langle \Psi | \hat{n}_1(\mathbf{r}', \mathbf{r}) | \Psi \rangle$.

We already know that the 1RDM enables to compute the electron density. Is it possible to restore the 1RDM from the electron density alone ?

Hint: show that the **full** one-electron **density matrix** is **required** for constructing **each elements** of the 1RDM.

Pair density operator

- Two-electron repulsion operator in the **r representation**:

$$\hat{E}_{pq} \longrightarrow \sum_{\sigma} \hat{\Psi}^{\dagger}(\mathbf{r}_1, \sigma) \hat{\Psi}(\mathbf{r}_2, \sigma)$$

$$\langle pr|qs\rangle \longrightarrow \langle \mathbf{r}_1 \mathbf{r}_2 | \hat{w}_{ee} | \mathbf{r}_3 \mathbf{r}_4 \rangle = \frac{1}{|\mathbf{r}_1 - \mathbf{r}_2|} \delta(\mathbf{r}_1 - \mathbf{r}_3) \delta(\mathbf{r}_2 - \mathbf{r}_4)$$

$$\hat{W}_{ee} = \frac{1}{2} \sum_{p,q,r,s} \langle pr|qs\rangle \left(\hat{E}_{pq} \hat{E}_{rs} - \delta_{qr} \hat{E}_{ps} \right) \longrightarrow \boxed{\hat{W}_{ee} = \frac{1}{2} \int \int d\mathbf{r}_1 d\mathbf{r}_2 \frac{\hat{n}_2(\mathbf{r}_1, \mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|}}$$

where $\hat{n}_2(\mathbf{r}_1, \mathbf{r}_2) = \hat{n}(\mathbf{r}_1) \hat{n}(\mathbf{r}_2) - \delta(\mathbf{r}_1 - \mathbf{r}_2) \hat{n}(\mathbf{r}_1)$ ← pair density operator !

Pair density operator

- **Expectation value** for the two-electron repulsion:

$$\langle \Psi | \hat{W}_{ee} | \Psi \rangle = \frac{1}{2} \int \int d\mathbf{r}_1 d\mathbf{r}_2 \frac{1}{|\mathbf{r}_1 - \mathbf{r}_2|} \left(\langle \Psi | \hat{n}(\mathbf{r}_1) \hat{n}(\mathbf{r}_2) | \Psi \rangle - \delta(\mathbf{r}_1 - \mathbf{r}_2) n_{\Psi}(\mathbf{r}_1) \right)$$

- If Ψ is the ground-state wavefunction of an interacting electronic system ($\hat{w} \neq 0$), it is usual in **density-functional theory** (DFT) to consider the **Kohn-Sham** (KS) Slater **determinant** Φ^{KS} that is the ground-state wavefunction of the non-interacting system ($\hat{w} = 0$) that has exactly the **same electron density**:

$$n_{\Phi^{KS}}(\mathbf{r}_1) = n_{\Psi}(\mathbf{r}_1),$$

thus leading to

$$\langle \Psi | \hat{W}_{ee} | \Psi \rangle - \langle \Phi^{KS} | \hat{W}_{ee} | \Phi^{KS} \rangle = \frac{1}{2} \int \int d\mathbf{r}_1 d\mathbf{r}_2 \frac{1}{|\mathbf{r}_1 - \mathbf{r}_2|} \left[\langle \Psi | \hat{n}(\mathbf{r}_1) \hat{n}(\mathbf{r}_2) | \Psi \rangle - \langle \Phi^{KS} | \hat{n}(\mathbf{r}_1) \hat{n}(\mathbf{r}_2) | \Phi^{KS} \rangle \right]$$

- This expression can be used in the calculation of the **correlation energy** (adiabatic connection fluctuation-dissipation theorem)

Hubbard Hamiltonian

$$h_{ij} \longrightarrow -t(\delta_{i,j-1} + \delta_{i,j+1}) + v_i \delta_{ij}$$

$$\langle ij|kl\rangle \longrightarrow U\delta_{ij}\delta_{ik}\delta_{lj}$$

$$\hat{E}_{ik}\hat{E}_{jl} - \delta_{kj}\hat{E}_{il} \longrightarrow \hat{n}_i\hat{n}_i - \hat{n}_i$$

where $\hat{n}_i = \hat{E}_{ii} = \hat{n}_{i\uparrow} + \hat{n}_{i\downarrow}$ so that $\hat{n}_i\hat{n}_i = 2\hat{n}_{i\uparrow}\hat{n}_{i\downarrow} + \hat{n}_i$

$$\hat{H} \longrightarrow \underbrace{-t \sum_{\langle i,j \rangle} \sum_{\sigma=\uparrow,\downarrow} \hat{a}_{i,\sigma}^\dagger \hat{a}_{j,\sigma}}_{\hat{\mathcal{T}} \text{ (hopping)}} + \underbrace{U \sum_i \hat{n}_{i\uparrow} \hat{n}_{i\downarrow}}_{\text{on-site repulsion}} + \underbrace{\sum_i v_i \hat{n}_i}_{\text{local potential}}$$

Spin operators

- **Spin vector operator** in second quantization:

$$\hat{\mathbf{S}} = \sum_{i=1}^N \hat{\mathbf{s}}(i) \equiv \sum_{pq, \sigma \sigma'} \underbrace{\langle \phi_p, \sigma | \hat{\mathbf{S}} | \phi_q, \sigma' \rangle}_{\langle \phi_p, \sigma | \phi_q, \hat{\mathbf{S}} | \sigma' \rangle} \hat{a}_{p, \sigma}^\dagger \hat{a}_{q, \sigma'}$$

$$\langle \phi_p, \sigma | \phi_q, \hat{\mathbf{S}} | \sigma' \rangle = \delta_{pq} \langle \sigma | \hat{\mathbf{S}} | \sigma' \rangle = \delta_{pq} \mathbf{s}_{\sigma \sigma'}$$

$$\text{with } \hat{s}^x \equiv \frac{1}{2} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad \hat{s}^y \equiv \frac{1}{2} \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad \hat{s}^z \equiv \frac{1}{2} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix},$$

thus leading to

$$\boxed{\hat{\mathbf{S}} \equiv \sum_p \hat{\mathbf{S}}_p} \quad \text{where}$$

$$\hat{S}_p^x = \frac{1}{2} \left(\hat{a}_{p, \uparrow}^\dagger \hat{a}_{p, \downarrow} + \hat{a}_{p, \downarrow}^\dagger \hat{a}_{p, \uparrow} \right), \quad \hat{S}_p^y = \frac{1}{2i} \left(\hat{a}_{p, \uparrow}^\dagger \hat{a}_{p, \downarrow} - \hat{a}_{p, \downarrow}^\dagger \hat{a}_{p, \uparrow} \right), \quad \hat{S}_p^z = \frac{1}{2} \left(\hat{a}_{p, \uparrow}^\dagger \hat{a}_{p, \uparrow} - \hat{a}_{p, \downarrow}^\dagger \hat{a}_{p, \downarrow} \right).$$

EXERCISE:

(1) Show that
$$[\hat{A}, \hat{B}\hat{C}] = [\hat{A}, \hat{B}] \hat{C} + \hat{B} [\hat{A}, \hat{C}] = [\hat{A}, \hat{B}]_+ \hat{C} - \hat{B} [\hat{A}, \hat{C}]_+.$$

(2) Deduce from (1) that
$$[\hat{a}_{p,\sigma}^\dagger \hat{a}_{q,\sigma'}, \hat{a}_{r,\tau}^\dagger \hat{a}_{s,\tau'}] = \delta_{rq} \delta_{\tau\sigma'} \hat{a}_{p,\sigma}^\dagger \hat{a}_{s,\tau'} - \delta_{sp} \delta_{\tau'\sigma} \hat{a}_{r,\tau}^\dagger \hat{a}_{q,\sigma'}.$$

We suggest to use in the following the notation
$$\hat{S}_p = \sum_{\sigma\sigma'} s_{\sigma\sigma'} \hat{a}_{p,\sigma}^\dagger \hat{a}_{p,\sigma'}.$$

(3) Explain why
$$[\hat{S}^x, \hat{S}^y] = \sum_p [\hat{S}_p^x, \hat{S}_p^y].$$

(4) Prove that
$$[\hat{S}_p^x, \hat{S}_p^y] = i \hat{S}_p^z.$$
 Hint: use (2) and the equality $[\hat{s}^x, \hat{s}^y] = i \hat{s}^z.$

(5) Conclude that \hat{S} is an **angular momentum** operator:
$$[\hat{S}^x, \hat{S}^y] = i \hat{S}^z.$$

Spin operators

- Another important model Hamiltonian is the **Heisenberg** Hamiltonian:

$$\hat{H} \longrightarrow -J \sum_p \hat{\mathbf{S}}_p \cdot \hat{\mathbf{S}}_{p+1}$$